

Table 8.1.1: Distribution of Pre dialysis Systolic Blood Pressure (mmHg), HD patients 1997-2004

Year	No. of subjects	Mean	SD	Median	LQ	UQ	% patients <120 mmHg	% patients 120- <140 mmHg	% patients 140- <160 mmHg	% patients 160- <180 mmHg	% patients ≥180 mmHg
1997	1659	144.5	20.8	144.2	130.8	158.1	11	30	35	19	4
1998	2108	146	20.5	146.7	133.2	159.2	10	27	39	19	5
1999	2965	148.7	20.8	148.5	135.3	162.2	8	25	38	23	6
2000	4310	148	20.6	147.8	134.8	161.7	9	25	38	23	6
2001	5147	148.8	20.9	148.8	134.9	162.6	8	25	37	23	7
2002	5911	149.2	20.6	149	135.8	163.3	8	24	38	24	6
2003	6839	149.7	20.2	149.8	136.4	162.9	7	24	39	23	7
2004	7856	149.8	20	150	136.7	163.2	7	23	39	25	6

Figure 8.1.1: Cumulative distribution of Pre dialysis Systolic Blood Pressure (mmHg), HD patients 1997-2004




Table 8.1.2: Distribution of Pre dialysis Systolic Blood Pressure (mmHg), CAPD patients 1997-2004

Year	No. of subjects	Mean	SD	Median	LQ	UQ	% patients <120 mmHg	% patients 120- <140 mmHg	% patients 140- <160 mmHg	% patients 160- <180 mmHg	% patients ≥180 mmHg
1997	468	142.7	20.3	142.9	128.3	156.3	13	31	37	17	3
1998	519	141	21.2	140	126.4	157.5	16	34	29	18	3
1999	576	141	19.8	140	127.2	156	14	35	34	15	2
2000	638	137.2	20.4	136.1	123.3	150	18	39	29	13	2
2001	739	139	20.2	137.5	125.8	151.7	16	38	30	13	3
2002	843	139.8	20.5	140	127.1	151.8	14	36	34	12	4
2003	1156	140.5	20.1	140	126.7	154.1	15	35	32	15	3
2004	1260	141	19.8	140.9	127.5	154.4	13	34	35	13	3

Figure 8.1.2: Cumulative distribution of Pre dialysis Systolic Blood Pressure (mmHg), CAPD patients 1997-2004




Table 8.1.3: Distribution of Pre dialysis Diastolic Blood Pressure (mmHg), HD patients 1997-2004

Year	No. of subjects	Mean	SD	Median	LQ	UQ	% patients <70 mmHg	% patients 70-<80 mmHg	% patients 80-<90 mmHg	% patients 90-<100 mmHg	% patients ≥100 mmHg
1997	1660	83.7	10.9	84.2	77	90.7	10	23	38	22	6
1998	2108	83.5	10.7	83.9	76.9	90.6	10	24	38	23	5
1999	2965	83.5	10.5	83.5	77.1	90	10	24	40	21	6
2000	4309	82.2	10.4	82.3	75.7	89	11	28	39	18	4
2001	5146	81.6	10.4	81.7	75	88.3	12	30	37	17	4
2002	5907	81.2	10.4	81.3	74.5	88.1	13	30	37	16	3
2003	6837	80.6	10.2	80.8	73.9	87.2	14	32	37	14	3
2004	7854	80.3	10.2	80.4	73.6	87	15	32	36	14	3

Figure 8.1.3: Cumulative distribution of Pre dialysis Diastolic Blood Pressure (mmHg), HD patients 1997-2004




Table 8.1.4: Distribution of Pre dialysis Diastolic Blood Pressure (mmHg), CAPD patients 1997-2004

Year	No. of subjects	Mean	SD	Median	LQ	UQ	% patients <70 mmHg	% patients 70-<80 mmHg	% patients 80-<90 mmHg	% patients 90-<100 mmHg	% patients ≥100 mmHg
1997	467	85.3	10.6	85.8	79.8	91.4	6	19	41	26	8
1998	519	84.3	11.3	85	77.1	90.1	8	24	36	24	8
1999	576	84	10.9	84.2	77.9	90	9	20	44	20	7
2000	638	82.9	11	83.3	76.6	89.6	10	24	41	20	5
2001	739	83.1	10.9	82.7	76.4	89.6	9	29	38	18	6
2002	843	82.8	10.8	83.4	76.1	90	11	24	41	21	5
2003	1158	82.2	10.9	82.3	75.5	89.4	12	26	38	19	4
2004	1259	82.2	10.5	83	75.4	89.2	11	28	39	18	4

Figure 8.1.4: Cumulative distribution of Pre dialysis Diastolic Blood Pressure (mmHg), CAPD patients 1997-2004




Table 8.1.5(a): Variation in BP control among HD centres 2004

(a) Median Systolic blood pressure (mmHg) among HD patients

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	46	120.5	130.4	140	144.9	151.9	158	159.8
1998	46	128.7	134.8	141.3	146.8	150	157.7	158.1
1999	69	132.1	135.5	143	149	155.3	164.5	170.2
2000	99	130.6	135	141.2	147.5	154.3	163	173.8
2001	116	126.7	136.3	143.1	148.8	155.7	162.9	185.8
2002	133	122.9	137.6	144.3	149	153.8	163.2	171.4
2003	156	124.6	136.3	144.5	150.8	155.6	163.2	171.4
2004	184	122.1	136.9	144.7	150.1	154.7	162.9	171

Figure 8.1.5(a): Variation in median systolic blood pressure (mmHg) among HD patients, HD centres 2004




Table 8.2.3: Distribution of serum Triglyceride (mmol/L), HD patients 1997-2004

Year	No. of subjects	Mean	SD	Median	LQ	UQ	% patients <1.7 mmol/L	% patients 1.7-<2.3 mmol/L	% patients 2.3-<3.5 mmol/L	% patients ≥3.5 mmol/L
1997	1074	2.1	1.4	1.8	1.3	2.5	45	24	18	12
1998	1089	2.2	1.5	1.8	1.3	2.6	42	26	20	12
1999	1634	2.1	1.3	1.7	1.2	2.5	49	22	18	11
2000	2393	2.1	1.4	1.7	1.3	2.6	48	22	19	12
2001	3162	2.1	1.4	1.7	1.2	2.5	48	22	17	13
2002	3861	2.1	1.4	1.8	1.2	2.5	47	22	18	12
2003	4715	2	1.3	1.7	1.2	2.5	48	23	18	11
2004	5543	2	1.2	1.7	1.2	2.4	51	22	17	10

Figure 8.2.3: Cumulative distribution of serum Triglyceride (mmol/L), HD patients 1997-2004




Table 8.2.4: Distribution of serum Triglyceride (mmol/L), CAPD patients 1997-2004

Year	No. of subjects	Mean	SD	Median	LQ	UQ	% patients <1.7 mmol/L	% patients 1.7-<2.3 mmol/L	% patients 2.3-<3.5 mmol/L	% patients ≥3.5 mmol/L
1997	413	2.6	1.9	2.2	1.4	3	36	22	25	18
1998	344	2.4	1.8	1.8	1.3	3	42	22	17	19
1999	421	2.4	1.6	2	1.4	3	38	25	18	19
2000	520	2.7	2.2	2.1	1.5	3	33	24	23	21
2001	576	2.6	1.8	2	1.4	3	36	22	22	20
2002	767	2.5	1.7	2	1.4	3	39	21	22	18
2003	1102	2.3	1.6	1.8	1.3	2.8	45	20	21	14
2004	1224	2.2	1.6	1.8	1.3	2.6	47	22	17	13

Figure 8.2.4: Cumulative distribution of serum Triglyceride (mmol/L), CAPD patients 1997-2004




Table 8.2.5(a): Variation in dyslipidaemia among HD centres 2004

(a) Median serum cholesterol level among HD patients

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	34	4.1	4.3	4.6	5	5.3	5.8	5.9
1998	28	4	4.4	4.8	5	5.2	5.3	5.4
1999	47	3.5	4.3	4.6	4.8	5.1	5.5	5.8
2000	75	4	4.3	4.8	4.9	5.2	5.5	5.8
2001	95	4.2	4.5	4.7	4.9	5.2	5.8	6.3
2002	112	4.4	4.5	4.7	4.9	5.1	5.4	6
2003	142	4	4.3	4.6	4.8	5	5.3	5.6
2004	164	3.8	4.1	4.5	4.7	4.9	5.3	5.7

Figure 8.2.5(a): Variation in median serum cholesterol level among HD patients, HD centres 2004




Table 8.2.5(b): Variation in dyslipidaemia among HD centres 2004

(b) Proportion of patients with serum cholesterol < 5.3 mmol/L

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	34	33	35	48	57	64	84	92
1998	28	38	42	52	59.5	69	84	89
1999	47	34	38	57	68	76	84	93
2000	75	32	38	51	60	68	83	100
2001	95	13	37	54	60	68	80	90
2002	112	31	42	57	64	70	79	92
2003	142	40	48	61	68	76	84	100
2004	164	39	50	63.5	71	78	89	96

Figure 8.2.5(b): Variation in proportion of patients with serum cholesterol < 5.3 mmol/L, HD centres 2004




Table 8.2.5(c): Variation in dyslipidaemia among HD centres 2004

(c) Median serum triglyceride level among HD patients

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	33	1.3	1.3	1.6	1.8	1.9	2.5	2.9
1998	27	1.4	1.5	1.7	1.8	1.9	2	2.2
1999	39	1.3	1.3	1.5	1.6	1.9	2.3	2.5
2000	59	1.3	1.4	1.5	1.8	2	2.4	2.6
2001	79	.9	1.3	1.5	1.7	1.9	2.5	3.3
2002	95	1.2	1.3	1.6	1.8	1.9	2.3	2.9
2003	122	1.2	1.3	1.6	1.7	1.9	2.2	2.5
2004	150	1	1.3	1.5	1.6	1.9	2.2	2.8

Figure 8.2.5(c): Variation in median serum triglyceride level among HD patients, HD centres 2004




Table 8.2.5(d): Variation in dyslipidaemia among HD centres 2004

(d) Proportion of patients with serum triglyceride < 2.1 mmol/L

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	33	29	36	60	65	77	83	85
1998	27	47	53	58	64	69	79	82
1999	39	43	48	61	67	75	90	92
2000	59	17	36	57	65	72	84	100
2001	79	33	41	57	66	74	86	100
2002	95	9	45	58	65	72	83	90
2003	122	30	48	59	67	74	86	100
2004	150	27	46	60	69	77	86	100

Figure 8.2.5(d): Variation in proportion of patients with serum triglyceride < 2.1 mmol/L, HD centres 2004




Table 8.2.6(a): Variation in dyslipidaemia among CAPD centres 2004

(a) Median serum cholesterol level among CAPD patients

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	6	5.8	5.8	5.9	5.9	6.1	6.1	6.1
1998	6	5.1	5.1	5.6	5.8	6.1	6.2	6.2
1999	8	5.3	5.3	5.6	5.7	5.9	6	6
2000	10	4.9	4.9	5.3	5.7	6	6.4	6.4
2001	10	5	5	5.5	5.9	6	6.2	6.2
2002	14	4.9	4.9	5.4	5.6	5.7	6.3	6.3
2003	17	4.6	4.6	5.2	5.3	5.7	6.1	6.1
2004	17	4.6	4.6	5	5.3	5.5	5.9	5.9

Figure 8.2.6(a): Variation in median serum cholesterol level among CAPD patients, CAPD centres 2004




Table 8.2.6(b): Variation in dyslipidaemia among CAPD centres 2004

(b) Proportion of patients with serum cholesterol < 5.3 mmol/L

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	6	27	27	27	29	31	33	33
1998	6	24	24	27	32	38	50	50
1999	8	6	6	29.5	39.5	43.5	50	50
2000	10	11	11	18	31	47	50	50
2001	10	23	23	30	34	44	61	61
2002	14	25	25	36	41	43	80	80
2003	17	0	0	38	48	54	81	81
2004	17	34	34	44	52	58	69	69

Figure 8.2.6(b): Variation in proportion of patients with serum cholesterol < 5.3 mmol/L, CAPD centres 2004




Table 8.2.6(c): Variation in dyslipidaemia among CAPD centres 2004

(c) Median serum triglyceride level among CAPD patients

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	6	1.7	1.7	1.9	2.1	2.2	2.4	2.4
1998	6	1.2	1.2	1.5	1.7	1.9	2.1	2.1
1999	8	1.6	1.6	1.9	2	2.1	2.6	2.6
2000	10	1.8	1.8	2	2.1	2.5	2.8	2.8
2001	10	1.5	1.5	1.8	2	2.1	3	3
2002	14	1.6	1.6	1.8	2	2.1	2.4	2.4
2003	17	1.1	1.1	1.7	1.8	2	2.2	2.2
2004	17	1.4	1.4	1.6	1.8	1.9	2.1	2.1

Figure 8.2.6(c): Variation in median serum triglyceride level among CAPD patients, CAPD centres 2004




Table 8.2.6(d): Variation in dyslipidaemia among CAPD centres 2004

(d) Proportion of patients with serum triglyceride < 2.1 mmol/L

Year	No. of centres	Min	5 th Centile	LQ	Median	UQ	95 th Centile	Max
1997	6	40	40	46	52	56	61	61
1998	6	51	51	58	60.5	70	73	73
1999	8	37	37	53	56.5	62.5	71	71
2000	10	18	18	38	51.5	54	67	67
2001	10	28	28	51	52.5	58	68	68
2002	14	37	37	51	54	57	75	75
2003	17	48	48	56	59	62	100	100
2004	17	49	49	60	62	65	93	93

Figure 8.2.6(d): Variation in proportion of patients with serum triglyceride < 2.1 mmol/L, CAPD centres 2004

